






Oxivir[®] Tb Hospital Grade One-Step Cleaner Disinfectant

| | | |
|--|--|--|
|  | <ol style="list-style-type: none"> 1. Apply Oxivir Tb undiluted to surfaces by spray, cloth or disposable wipe. | <p>EPA Reg. 70627-56</p> <p>Tested & proven effective against a wide spectrum of pathogens:</p> <p>Viruses:</p> <ul style="list-style-type: none"> • Hepatitis B • Hepatitis C • Herpes Simplex Virus, Type 1 & 2 • HIV-1 (AIDS virus) • Human Coronavirus • Influenza A • Norovirus • Poliovirus • Rotavirus • Adenovirus • Rhinovirus Type 37 • Avian influenza A • Feline Calicivirus <p>Bacteria:</p> <ul style="list-style-type: none"> • Pseudomonas aeruginosa • Salmonella enterica • Staphylococcus aureus • Acinetobacter baumannii • Escherichia coli O157:H7 • Klebsiella pneumoniae • Escherichia coli with extended beta-lactamase resistance (ESBL) • Shigella Dysenteriae <p>Antibiotic- Resistant Bacteria:</p> <ul style="list-style-type: none"> • Enterococcus faecalis (VRE) • Staphylococcus aureus (MRSA) • Staphylococcus aureus (CA-MRSA) • Staphylococcus aureus (CA-MRSA) <p>Fungi: Trichophyton mentagrophytes</p> <p>Tb: Mycobacterium bovis</p> |
|  | <ol style="list-style-type: none"> 1. All surfaces must remain visibly wet for 1 minute for viruses and bacteria. Allow 5 minutes for Tb and 10 minutes for Fungi. 2. Allow to air dry. 3. If concerned about streaking, wipe with a microfibre cloth. | |
|  | <ol style="list-style-type: none"> 1. If disinfecting food contact surfaces, rinse with water 2. If Clostridium Difficile outbreak is identified, use microfibre . Or use Sporicide Plus. | |
| <p><u>SUPPLIES/TOOLS:</u></p> <ul style="list-style-type: none"> • Ultra-microfibre cloths • Disposable dry towel • Gloves should be considered, as prolonged hand contact will dry skin | <p><u>SAFETY:</u></p> <p>No PPE Gear Needed.</p> <ul style="list-style-type: none"> • Oxivir Tb has a low concentration of active (AHP) • Wear utility gloves and other PPE appropriate to the task such as site decontamination after spills of blood or other potentially infectious materials (NHMRC Guidelines). | |